

# Cancer Related Inhibitors

For more information : <http://www.funakoshi.co.jp/exports>

## Inhibitor for DNA Helicase


### Heliquinomycin

Heliquinomycin inhibits DNA helicase from HeLa cell in a non-competitive manner with the inhibition constant (Ki) of 6.8 mM. The topoisomerase II and I enzymes are inhibited at 30 µg/ml and 100 µg/ml of heliquinomycin, respectively. Heliquinomycin inhibits the growth of HeLa S3, KB, LS180, K562 and HL60 human tumor cell lines at IC<sub>50</sub> values of 0.96 to 2.8 µg/ml. Heliquinomycin inhibits both DNA and RNA synthesis in cell culture but does not inhibit protein synthesis. HeLa S3 cells are arrested at the G2/M phase by heliquinomycin.

This product is licensed under JP patent NO.3490154

#### Reference

- Chino, M., *et al.*, *J. Antibiotics*, 49, 752~757 (1996).  
Chino, M., *et al.*, *J. Antibiotics*, 50, 143~146 (1997).  
Chino, M., *et al.*, *J. Antibiotics*, 50, 781~784 (1997).  
Chino, M., *et al.*, *J. Antibiotics*, 51, 480~486 (1998).




[ Manufacturer : IMC ]

Product Name	Size	Catalog #	Storage
Heliquinomycin	0.2 mg	10665	4°C

## Inhibitor for Heparanase

### Heparastatin <SF4>

Heparastatin (SF4) inhibits recombinant human heparanase from human melanoma A375M cells transfected with pBK-CMV expression vectors containing the heparanase cDNA with IC<sub>50</sub> 1.02 µM. Heparastatin (SF4) inhibits β-D-glucuronidase from bovine liver with IC<sub>50</sub> 6.5 x 10<sup>-2</sup> µM. Heparastatin (SF4) (100 µM) completely inhibits the enzyme activity of recombinant heparanase of murine mammary epithelial cells (NMuMG) transfected with a mouse heparanase expression vector pcDNA3.1(-)-Hygro-Hep at 0.15 µg/ml in a *in vitro* HS degradation assay. Heparastatin (SF4) inhibits heparan sulfate (HS) chain degradation of HSPGs of Matrigel by heparanase of the LPS-treated microglial lysates from the forebrain cells of Wistar rats and the *in vitro* transmigration of microglia through the Matrigel-coated insert in a dose-dependent manner. Heparastatin (SF4) markedly inhibits degradation of HS by heparanase in the nucleus translocated from the cytoplasm of the calcium-induced human esophageal keratinocyte cells and keratinocyte differentiation at 100µM. Heparastatin (SF4) markedly inhibits in a dose-dependent manner experimentally induced pulmonary metastasis of the B16BL6 in mice. Inhibitory ratio by *ex vivo* treatment with 50 µg/ml of Heparastatin (SF4) is 90.8%. Heparastatin (SF4) shows 57% inhibition of lung metastasis of 3LL cells by s.c. inoculation in mice with i.v. administration of 100 mg/kg/day for 5 days.



[ Manufacturer : IMC ]

Product Name	Size	Catalog #	Storage
Heparastatin <SF4>	0.2 mg	11829	4°C

## Inhibitor for Protein Phosphatase 2A


### Cytostatin

Cytostatin inhibits the adhesion of B16 melanoma cells to laminin and collagen type IV in a dose dependent manner but not to fibronectin. The IC<sub>50</sub> values are 1.3 µg/ml to laminin and 1.4 µg/ml to collagen. The administration of cytostatin inhibits metastases of B16-F10 markedly. The inhibitory ration is about 60 to 70% at 1.25 mg/kg/day. Cytostatin inhibits protein phosphatase 2A with an IC<sub>50</sub> of 0.09 µg/ml in a non competitive manner against a substrate, p-nitrophenyl phosphate, but it has no apparent effect on other protein phosphatases including protein phosphatase 1, protein phosphatase 2B and alkaline phosphatase even at 100 µg/ml.

This product is licensed under JP patent NO.367455

#### Reference

- Amemiya, M., *et al.*, *J. Antibiotics*, 47, 536~540 (1994).  
 Masuda, T., *et al.*, *J. Antibiotics*, 48, 528~529 (1995).  
 Kawada, M., *et al.*, *Biochim. Biophys. Acta.*, 1452, 209~217 (1999).  
 Kawada, M., *et al.*, *Int. Immunopharmacol.*, 3, 179~188 (2003).



[ Manufacturer : IMC ]


Product Name	Size	Catalog #	Storage
Cytostatin	0.2 mg	10664	-20°C

### Rubratoxin A

Rubratoxin A is a mycotoxin produced by *Penicillium*. While this compound inhibits protein phosphatase 2A (PP2A) in a competitive manner with the inhibition constant (K<sub>i</sub>) of 28.7 nM (IC<sub>50</sub>= 6.1-15.1 nM), it does not inhibit protein phosphatase 1, protein phosphatase 2B, protein tyrosine phosphatase 1B, alkaline phosphatase, and calf intestine phosphatase(CIP) up to 200 µM. Rubratoxin A shows cytotoxicity in cultured mammalian cells with IC<sub>50</sub> values of around 15 µM and induces overphosphorylation of PP2A substrate proteins.

#### Reference

- Wada, S., *et al.*, *Cancer Sci.*, 101, 743~750 (2010).



[ Manufacturer : IMC ]

Product Name	Size	Catalog #	Storage
Rubratoxin A	0.2 mg	10663	4°C

#### NOTE

- ※ All products here are research use only, not for diagnostic use.
- ※ Specs might be changed for improvement without notice.

- ※ Company name and product name are trademark or registered mark.
- ※ Please contact your local distributors for orders, quote request and inquiry.

Your Local Distributor



**Bio-REV Pte. Ltd.**

36 Toh Guan Road East, #01-39  
 Enterprise Hub, Singapore 608 580  
 Tel: (65) 6273-3022  
 Fax: (65) 6273-3020  
 Email: sales@bio-rev.com  
 Technical Support: techserv@bio-rev.com



**funakoshi Co., Ltd.**

Address: 9-7 Hongo 2-Chome, Bunkyo-ku,  
 Tokyo 113-0033 JAPAN  
 Phone : +81-3-5684-6296  
 Fax : +81-3-5684-6297  
 Email : export@funakoshi.co.jp